


Medication Orders and Anaphylaxis Action Plan

For those requiring emergency EPINEPHRINE treatment
 "Anaphylaxis is a serious allergic reaction that is rapid in onset and may cause death."
 (National Institute of Allergy & Infectious Disease, 2010)



Name/DOB: _____

ALLERGIC to: _____

History of Asthma: Yes (more at risk for severe reaction) No

May self-carry medications: Yes No

May self administer medications: Yes No

Medication Doses

EPINEPHRINE Dose:

Up to 55 lbs. (25kg)

- EpiPen Jr. (0.15mg)
- Adrenaclick (0.15mg)
- Auvi-Q (0.15mg)

Over 55 lbs. (25kg)

- EpiPen (0.3mg)
- Adrenaclick (0.3mg)
- Auvi-Q (0.3mg)

***Antihistamine Type + Dose:**

- Benadryl (also known as Diphenhydramine)
 - 12.5 mg (1 teaspoon or 1 chewable)
 - 25 mg (2 teaspoons or 2 chewables)
 - 50 mg (4 teaspoons or 4 chewables)
- Other antihistamine: _____

Extremely reactive to the following foods: _____

THEREFORE:

- If checked, give EPINEPHRINE immediately for ANY symptoms if the allergen was *likely* eaten.
- If checked, give EPINEPHRINE immediately if the allergen was *definitely* eaten, even if no symptoms are noted.


Any SEVERE SYMPTOMS after suspected or known ingestion:

One or more of the following:

- Lung: Short of breath, wheeze, repetitive cough
- Heart: Pale, blue, faint, weak pulse, dizzy, confused
- Throat: Tight, hoarse, trouble breathing/swallowing
- Mouth: Obstructive swelling (tongue and/or lips)
- Skin: Many hives over body
- Other: Anxiety, confusion, feeling something bad is about to happen

Or **combination** of symptoms from different body areas:

- Skin: Hives, itchy rashes, swelling (eyes, lips)
- Gut: Vomiting, crampy pain



1. INJECT EPINEPHRINE


IMMEDIATELY

2. Call 911
3. Begin monitoring (as specified below)
4. Give additional medications:*
 - Antihistamine
 - Inhaler (bronchodilator) if asthma

*Antihistamines & inhalers/bronchodilators are not to be depended upon to treat severe reaction (anaphylaxis). USE EPINEPHRINE

MILD SYMPTOMS only:

- Mouth: Itchy Mouth
- Skin: A few hives around mouth/face, mild itch
- Gut: Mild nausea/discomfort



1. GIVE ANTIHISTAMINE

2. Stay with student; alert healthcare professionals and parent/guardian
3. If symptoms progress (see above) USE EPINEPHRINE
4. Begin monitoring (as specified below)

For unique situations: _____

Monitoring

A **SECOND DOSE** of EPINEPHRINE can be given 5 minutes or more after the first if symptoms persist or recur.

Stay with person; alert healthcare professionals and parent/guardian. Tell rescue squad EPINEPHRINE was given. Note time when EPINEPHRINE was administered. For a severe reaction, consider keeping person lying on back with legs raised. Treat person even if parents cannot be reached. Send used epinephrine injector with EMS. See back/attached for auto-injection technique.

Provider Signature: _____ Phone _____ Date _____

Printed Name: _____

Parent/Guardian Signature: _____ Phone _____ Date _____



An affiliate of Children's Hospitals and Clinics of Minnesota
 www.clinics4kids.org


Turn Form Over →

EpiPEN® Auto-Injector and EpiPEN Jr® Auto-Injector Directions

- First, remove the EPI PEN Auto-Injector from the plastic carrying case



- Pull off the blue safety release cap
- Hold orange tip near outer thigh (always apply to thigh)




- Swing and firmly push orange tip against outer thigh. Hold on thigh for approximately 10 seconds.
- Remove the EPIPEN Auto-Injector and massage the area for 10 more seconds




EpiPen®, EpiPen 2-Pak®, and EpiPen Jr 2-Pak® are registered trademarks of Mylan Inc. licensed exclusively to its wholly-owned subsidiary, Mylan Specialty LP.

Auvi-Q® 0.3 mg and Auvi-Q® 0.15 mg Directions

1. Remove the outer case of Auvi-Q. This will automatically activate the voice instructions.
2. Pull off RED safety guard.




3. Place black end against outer thigh, then press firmly and hold for 5 seconds.




©2002-2013 sanofi-aventis U.S. LLC. All rights reserved

Adrenaclick® 0.3 mg and Adrenaclick® 0.15 mg Directions



Remove GREY caps labeled "1" and "2".

Place RED rounded tip against Outer thigh, press down hard until needle penetrates. Hold for 10 seconds, then remove.



A food allergy response kit should contain at least two doses of epinephrine, other medications as noted by the student's physician, and a copy of this Food Allergy Action Plan. A kit must accompany the student if he/she is off school grounds (i.e., fieldtrip).

Contacts

Call 911 (Rescue squad: _____) Doctor: _____ Phone: _____

Parent/Guardian: _____ Phone: _____

Other Emergency Contacts

Name/Relationship: _____ Phone: _____

Name/Relationship: _____ Phone: _____



An affiliate of Children's Hospitals and Clinics of Minnesota
www.clinics4kids.org

Adapted from the Food Allergy & Anaphylaxis Network (FAAN) Action Plan

www.foodallergy.org